INDEX TO VOLUME 24

Subject Index

Abortion

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Abortion reform in South Africa: a case study of the 1996 Choice on Termination of Pregnancy Act, Guttmacher S, Kapadia F, Naude JTW and de Pinho H, Special Report, p. 191.

Chilean women prosecuted after clandestine abortion often are reported by the hospitals that treat them, see **Latin America and the Caribbean**.

Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, see **Fertility: Planning and Effects.**

Global review of laws on induced abortion, 1985–1997, Rahman A, Katzive L and Henshaw SK, p. 56.

Incidence of induced abortion in Nigeria, see **Africa**.

Induced abortion in Matlab, Bangladesh: trends and determinants, Ahmed MK, Rahman M and van Ginneken J, p. 128.

Needed: access to contraception, see **Contraceptive Practice**.

Postabortion caseload in Egyptian hospitals: a descriptive study, Huntington D, Nawar L, Hassan ED, Youssef H and Abdel-Tawab N, p. 25.

Potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, Trinh HV, Bishop A, Vuong TH, Luong XH, Tran DC and Nguyen TI, p. 165.

Proactive providers, Update, p. 154.

"Quasi-legal" abortion services in a sub-Saharan setting: users' profile and motivations, Agadjanian V, p. 111.

Reasons why women have induced abortions: evidence from 27 countries, Bankole A, Singh S and Haas T, p. 117.

Africa

Abortion reform in South Africa: a case study of the 1996 Choice on Termination of Pregnancy Act, see **Abortion**.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, Digest, p. 46.

Consent: voluntary or coerced? see **Program Evaluation and Funding**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see Digest, p. 145.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Factors in HIV infectivity, see **Sexually Transmitted Diseases**.

Female circumcision is nearly universal in Egypt, Eritrea, Mali and Sudan, Digest, p. 47.

Global review of laws on induced abortion, see **Abortion**.

HIV saps breastfeeding benefits, see **Breastfeeding**.

Improved services for sexually transmitted diseases in Tanzania are cost-effective in cutting HIV rates, see **Sexually Transmitted Diseases**.

In Ghana and Kenya, many spouses disagree on their need for contraception, Digest, p. 100.

In Tanzania, ideal family size closely resembles actual number of children, Digest, p. 147.

Incidence of induced abortion in Nigeria, Henshaw SK, Singh S, Oye-Adeniran BA, Adewole IF, Iwere N and Cuca YP, p. 156.

Increasing access to family planning services in rural Mali through community-based distribution, Katz KR, West CG, Doumbia F and Kané F, p. 104.

Informed choice and decision-making in family planning counseling in Kenya, Kim YM, Kols A and Mucheke S, p. 4.

Maternal mortality in Africa, Update, p. 102.

Quality of family planning services for breastfeeding women in Senegal, Research Note, Stein K, Measham D and Winikoff B, p. 188.

"Quasi-legal" abortion services in a sub-Saharan setting: users' profile and motivations, see **Abortion**.

Rates of pregnancy and birth in rural Uganda are lower among HIV-infected women, Digest, p. 146.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Sexual activity and condom use in Lusaka, Zambia, Agha S, p. 32.

Sexual behavior and attitudes among unmarried urban youths in Guinea, Görgen R, Yansané ML, Marx M and Millimounou D, p. 65.

Teenage contraceptive needs in urban South Africa: a case study, Mfono Z, p. 180.

Time dynamics of unmet need: an example from Morocco, Westoff CF and Bankole A, p. 12.

Vaginal drying agents and HIV transmission, see **Sexual Behavior**.

When spouses disagree, see **Contraceptive Practice**.

Where have all the vaginal foaming tablets gone? program statistics and user dynamics in Ghana, Research Note, Steiner MJ, Attafuah JD, Stanback J and Nutley T, p. 91.

Asia

Among Uzbekistani women, little difference exists between actual and desired levels of fertility, Digest, p. 43.

As rise in family planning use slows in Bangladesh, method mix shifts toward reversible contraception, Digest, p. 201.

Challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, Comment, Hull TH, p. 176.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Family planning clinics through women's eyes and voices: a case study from rural Bangladesh, Schuler SR and Hossain Z, p. 170.

Family planning pays off, see **Fertility: Planning and Effects**.

Female sterilization in Nepal: a comparison of two types of service delivery, Thapa S and Friedman M, p. 78.

Global review of laws on induced abortion, see **Abortion**.

How education saves lives, see **Maternal and Child Health**.

In Pakistan, contraceptive prevalence is increasing, but fertility remains high, Digest, p. 203.

In Vietnam, small family size means better education among the young, but socioeconomic factors are key, Digest, p. 149.

India: how fast a decline? Update, p. 3.

Induced abortion in Matlab, Bangladesh: trends and determinants, see **Abortion**.

Involving men, helping women, see **Reproductive Health**.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, Digest, p. 95.

Major HIV risk factor facing many Thai women is the sexual behavior of their husband or partner, see **Sexual Behavior**.

Monogamous—and at risk of HIV, see **Sexually Transmitted Diseases**.

Poor socioeconomic status is linked to high maternal mortality in rural Pakistan, Digest, p. 96.

Potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, see **Abortion**.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Selective listening? see **Maternal and Child Health**.

Son preference in Anhui Province, China, Graham MJ, Larsen U and Xu X, p. 72.

Training by radio, Update, p. 54.

Two doses will do, see **Maternal and Child Health**.

INDEX TO VOLUME 24

Subject Index

Abortion

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Abortion reform in South Africa: a case study of the 1996 Choice on Termination of Pregnancy Act, Guttmacher S, Kapadia F, Naude JTW and de Pinho H, Special Report, p. 191.

Chilean women prosecuted after clandestine abortion often are reported by the hospitals that treat them, see **Latin America and the Caribbean**.

Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, see **Fertility: Planning and Effects.**

Global review of laws on induced abortion, 1985–1997, Rahman A, Katzive L and Henshaw SK, p. 56.

Incidence of induced abortion in Nigeria, see **Africa**.

Induced abortion in Matlab, Bangladesh: trends and determinants, Ahmed MK, Rahman M and van Ginneken J, p. 128.

Needed: access to contraception, see **Contraceptive Practice**.

Postabortion caseload in Egyptian hospitals: a descriptive study, Huntington D, Nawar L, Hassan ED, Youssef H and Abdel-Tawab N, p. 25.

Potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, Trinh HV, Bishop A, Vuong TH, Luong XH, Tran DC and Nguyen TI, p. 165.

Proactive providers, Update, p. 154.

"Quasi-legal" abortion services in a sub-Saharan setting: users' profile and motivations, Agadjanian V, p. 111.

Reasons why women have induced abortions: evidence from 27 countries, Bankole A, Singh S and Haas T, p. 117.

Africa

Abortion reform in South Africa: a case study of the 1996 Choice on Termination of Pregnancy Act, see **Abortion**.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, Digest, p. 46.

Consent: voluntary or coerced? see **Program Evaluation and Funding**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see Digest, p. 145.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Factors in HIV infectivity, see **Sexually Transmitted Diseases**.

Female circumcision is nearly universal in Egypt, Eritrea, Mali and Sudan, Digest, p. 47.

Global review of laws on induced abortion, see **Abortion**.

HIV saps breastfeeding benefits, see **Breastfeeding**.

Improved services for sexually transmitted diseases in Tanzania are cost-effective in cutting HIV rates, see **Sexually Transmitted Diseases**.

In Ghana and Kenya, many spouses disagree on their need for contraception, Digest, p. 100.

In Tanzania, ideal family size closely resembles actual number of children, Digest, p. 147.

Incidence of induced abortion in Nigeria, Henshaw SK, Singh S, Oye-Adeniran BA, Adewole IF, Iwere N and Cuca YP, p. 156.

Increasing access to family planning services in rural Mali through community-based distribution, Katz KR, West CG, Doumbia F and Kané F, p. 104.

Informed choice and decision-making in family planning counseling in Kenya, Kim YM, Kols A and Mucheke S, p. 4.

Maternal mortality in Africa, Update, p. 102.

Quality of family planning services for breastfeeding women in Senegal, Research Note, Stein K, Measham D and Winikoff B, p. 188.

"Quasi-legal" abortion services in a sub-Saharan setting: users' profile and motivations, see **Abortion**.

Rates of pregnancy and birth in rural Uganda are lower among HIV-infected women, Digest, p. 146.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Sexual activity and condom use in Lusaka, Zambia, Agha S, p. 32.

Sexual behavior and attitudes among unmarried urban youths in Guinea, Görgen R, Yansané ML, Marx M and Millimounou D, p. 65.

Teenage contraceptive needs in urban South Africa: a case study, Mfono Z, p. 180.

Time dynamics of unmet need: an example from Morocco, Westoff CF and Bankole A, p. 12.

Vaginal drying agents and HIV transmission, see **Sexual Behavior**.

When spouses disagree, see **Contraceptive Practice**.

Where have all the vaginal foaming tablets gone? program statistics and user dynamics in Ghana, Research Note, Steiner MJ, Attafuah JD, Stanback J and Nutley T, p. 91.

Asia

Among Uzbekistani women, little difference exists between actual and desired levels of fertility, Digest, p. 43.

As rise in family planning use slows in Bangladesh, method mix shifts toward reversible contraception, Digest, p. 201.

Challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, Comment, Hull TH, p. 176.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Family planning clinics through women's eyes and voices: a case study from rural Bangladesh, Schuler SR and Hossain Z, p. 170.

Family planning pays off, see **Fertility: Planning and Effects**.

Female sterilization in Nepal: a comparison of two types of service delivery, Thapa S and Friedman M, p. 78.

Global review of laws on induced abortion, see **Abortion**.

How education saves lives, see **Maternal and Child Health**.

In Pakistan, contraceptive prevalence is increasing, but fertility remains high, Digest, p. 203.

In Vietnam, small family size means better education among the young, but socioeconomic factors are key, Digest, p. 149.

India: how fast a decline? Update, p. 3.

Induced abortion in Matlab, Bangladesh: trends and determinants, see **Abortion**.

Involving men, helping women, see **Reproductive Health**.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, Digest, p. 95.

Major HIV risk factor facing many Thai women is the sexual behavior of their husband or partner, see **Sexual Behavior**.

Monogamous—and at risk of HIV, see **Sexually Transmitted Diseases**.

Poor socioeconomic status is linked to high maternal mortality in rural Pakistan, Digest, p. 96.

Potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, see **Abortion**.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Selective listening? see **Maternal and Child Health**.

Son preference in Anhui Province, China, Graham MJ, Larsen U and Xu X, p. 72.

Training by radio, Update, p. 54.

Two doses will do, see **Maternal and Child Health**.

Barrier Methods

Elevated condom breakage and slippage rates are linked to history of failure, specific types of misuse, Digest, p. 199.

Nonoxynol-9 fails to stop STDs, Update, p. 154.

Sexual activity and condom use in Lusaka, Zambia, see **Africa**.

Where have all the vaginal foaming tablets gone? program statistics and user dynamics in Ghana, see **Africa**.

Breastfeeding

Breastfeeding's HIV risk, Update, p. 155.

HIV saps breastfeeding benefits, Update, p. 3.

Lactational amenorrhea method provides effective postpartum contraception, Digest, p. 45.

Quality of family planning services for breastfeeding women in Senegal, see **Africa**.

Contraceptive Practice

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, see **Africa**.

As rise in family planning use slows in Bangladesh, method mix shifts toward reversible contraception, see **Asia**.

Catholicism and contraception, Update, p. 102.

Challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, see **Asia**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see **Africa**.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Elevated condom breakage and slippage rates are linked to specific types of misuse, see **Barrier Methods**.

Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, see **Fertility: Planning and Effects**.

Half of Paraguayan women use a method; almost eight in 10 births are planned, see **Latin America and the Caribbean**.

How to improve family planning and save lives using a stage-of-life approach, see **Population Policy and Research**.

In Ghana and Kenya, many spouses disagree on their need for contraception, see **Africa**.

In Pakistan, contraceptive prevalence is increasing, but fertility remains high, see **Asia**.

Increasing access to family planning services in rural Mali through community-based distribution, see **Africa**.

Lactational amenorrhea method provides effective postpartum contraception, see **Breastfeeding**.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, see **Asia**.

Needed: access to contraception, Update, p. 102.

Patterns of desired fertility and contraceptive use in Kuwait, see **Fertility: Planning and Effects**.

Teenage contraceptive needs in urban South Africa: a case study, see **Africa**.

Time dynamics of unmet need: an example from Morocco, see **Africa.**

When spouses disagree, Update, p. 2.

Europe

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, Digest, p. 150.

Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, see **Fertility: Planning and Effects**.

Needed: access to contraception, see **Contraceptive Practice**.

Fertility: Planning and Effects

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, see **Africa**.

Among Uzbekistani women, actual and desired fertility levels are similar, see **Asia**.

As rise in family planning use slows in Bangladesh, method mix shifts toward reversible contraception, see **Asia**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see **Africa**.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, Bankole A and Singh S, p. 15; and Correction, p. 138.

Family planning pays off, Update, p. 2.

Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, Research Note, Carlson E and Omori M, p. 184.

Half of Paraguayan women use a method; almost eight in 10 births are planned, see **Latin America and the Caribbean**.

In Pakistan, contraceptive prevalence is increasing, but fertility remains high, see **Asia.**

In Tanzania, ideal family size closely resembles actual number of children, see **Africa**.

In Turkey, women's fertility is linked to education, employment and freedom to choose a husband, see **Middle East**.

In Vietnam, small family size means better education among the young, but socioeconomic factors are key, see **Asia.**

India: how fast a decline? see **Asia**.

Lactational amenorrhea method provides effective postpartum contraception, see **Breastfeeding**.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, see **Asia**.

Patterns of desired fertility and contraceptive use in Kuwait, Shah NM, Shah MA and Radovanovic Z, p. 133.

Rates of pregnancy and birth in rural Uganda are lower among HIV-infected women, see **Africa**.

Son preference in Anhui Province, China, see **Asia**.

Time dynamics of unmet need: an example from Morocco, see **Africa**.

When spouses disagree, see **Contraceptive Practice**.

Hormonal Implants and Injectables

Challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, see **Asia**.

Injectable eases sickle-cell pain, Update, p. 54.

Intrauterine Devices

HIV-infected women who have access to health care may safely use IUD, Digest, p. 203.

12-year IUDs, Update, p. 55.

Latin America and the Caribbean

Chilean women prosecuted after clandestine abortion often are reported by the hospitals that treat them, Digest, p. 200.

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Gender differences in factors influencing first intercourse among urban students in Chile, see **Sexual Behavior**.

Global review of laws on induced abortion, see **Abortion**.

Half of Paraguayan women use a method; almost eight in 10 births are planned, Digest, p. 99.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Males

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

In Ghana and Kenya, many spouses disagree on their need for contraception, see **Africa.**

Involving men, helping women, see **Reproductive Health**.

Major HIV risk factor facing many Thai women is the sexual behavior of their husband or partner, see **Sexual Behavior**.

Men as partners in reproductive health: from issues to action, see **Reproductive Health**.

When spouses disagree, see **Contraceptive Practice**.

Maternal and Child Health

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, see **Africa**.

As rise in family planning use slows in Bangladesh, method mix shifts toward reversible contraception, see **Asia**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see **Africa**.

Half of Paraguayan women use a method; almost eight in 10 births are planned, see **Latin America and the Caribbean**.

How education saves lives, Update, p. 54.

How to improve family planning and save lives using a stage-of-life approach, see **Population Policy and Research**.

Lactational amenorrhea method provides effective postpartum contraception, see **Breastfeeding**.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, see **Asia**.

Maternal mortality in Africa, see **Africa**.

Mysteries of immunization, Update, p. 155.

Nourishment for two, Update, p. 2.

Poor socioeconomic status is linked to high maternal mortality in rural Pakistan, see **Asia**.

Selective listening? Update, p. 55.

Two doses will do, Update, p. 103.

Middle East

Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, see **Fertility: Planning and Effects**.

Global review of laws on induced abortion, see **Abortion**.

In Turkey, women's fertility is linked to education, employment and freedom to choose a husband, Digest, p. 97.

Patterns of desired fertility and contraceptive use in Kuwait, see **Fertility: Planning and Effects**.

Postabortion caseload in Egyptian hospitals: a descriptive study, see **Abortion**.

Reasons why women have induced abortions: evidence from 27 countries, see **Abortion**.

Oral Contraceptives

Quality of family planning services for breastfeeding women in Senegal, see **Africa**.

Population Policy and Research

Billions and billions of people, Update, p. 154.

Family planning pays off, see **Fertility: Planning and Effects**.

How to improve family planning and save lives using a stage-of-life approach, Viewpoint, Potts M, Rooks J and Holt BY, p. 195.

India: how fast a decline? see **Asia**.

Still growing, but more slowly, Update, p. 102.

Pregnancy and Childbearing

Café olé! Update, p. 154.

High-risk pregnant women, Update, p. 154.

HIV, vitamins and birth outcomes, Update, p. 155.

Program Evaluation and Funding

Challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, see **Asia**.

Consent: voluntary or coerced? Update, p. 103.

Family planning clinics through women's eyes and voices: a case study from rural Bangladesh, see **Asia**.

Female sterilization in Nepal: a comparison of two types of service delivery, see **Asia**.

Improved services for sexually transmitted diseases in Tanzania are cost-effective in cutting HIV rates, see **Sexually Transmitted Diseases**.

Increasing access to family planning services in rural Mali through community-based distribution, see **Africa**.

Informed choice and decision-making in family planning counseling in Kenya, see **Africa**.

Potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, see **Abortion**.

Quality of family planning services for breastfeeding women in Senegal, see **Africa**.

Teenage contraceptive needs in urban South Africa: a case study, see **Africa**.

Training by radio, see **Asia**.

What have we learned from studying changes in service guidelines and practices? Comment, Hardee K, Janowitz B, Stanback J and Villinski MT, p. 84.

Where have all the vaginal foaming tablets gone? program statistics and user dynamics in Ghana, see **Africa**.

Religious Attitudes

Catholicism and contraception, see **Contraceptive Practice**.

Reproductive Health

Do pollutants cause breast cancer? Update, p. 3.

Female circumcision is nearly universal in Egypt, Eritrea, Mali and Sudan, see **Africa**.

Involving men, helping women, Update, p. 54.

Men as partners in reproductive health: from issues to action, Special Report, Wegner MN Landry E, Wilkinson D and Tzanis J, p. 38

Sexual Behavior

Gender differences in factors influencing first intercourse among urban students in Chile, Murray NJ, Zabin LS, Toledo-Dreves V and Luengo-Charath X, p. 139.

Major HIV risk factor facing many Thai women is the sexual behavior of their husband or partner, Digest, p. 198.

Sex makes the world go 'round, Update, p. 103.

Sexual activity and condom use in Lusaka, Zambia, see **Africa**.

Sexual behavior and attitudes among unmarried urban youths in Guinea, see **Africa**.

Vaginal drying agents and HIV transmission, Viewpoint, Kun KE, p. 93.

Sexually Transmitted Diseases

Abortion rate among young Romanians declines; those not in union report rise in contraceptive use, see **Europe**.

Affordable HIV regimen, Update, p. 54.

Almost one in three Ugandan births are unwanted or mistimed, but few women practice contraception, see **Africa**.

Breastfeeding's HIV risk, see **Breastfeeding**.

Contraceptive knowledge is nearly universal in Senegal, yet fewer than one in seven women use a method, see **Africa**.

Cost-effective HIV prevention, Update, p. 102.

Factors in HIV infectivity, Update, p. 2.

HIV-infected women who have access to health care may safely use IUD, see **Intrauterine Devices**.

HIV prevalence revised upward, Update, p. 55.

HIV saps breastfeeding benefits, see **Breastfeeding**.

How to improve family planning and save lives using a stage-of-life approach, see **Population Policy and Research**.

Improved services for sexually transmitted diseases in Tanzania are cost-effective in cutting HIV rates, Digest, p. 98.

Levels of unwanted childbearing are high in Nepal, where only one in three women use contraceptives, see **Asia**.

Major HIV risk factor facing many Thai women is the sexual behavior of their husband or partner, see **Sexual Behavior**.

Monogamous—and at risk of HIV, Update, p. 2.

Nonoxynol-9 fails to stop STDs, see **Barrier Methods**.

Rates of pregnancy and birth in rural Uganda are lower among HIV-infected women, see **Africa**.

Vaginal drying agents and HIV transmission, see **Sexual Behavior**.

Status of Women

Female circumcision is nearly universal in Egypt, Eritrea, Mali and Sudan, see **Africa**.

How education saves lives, see **Maternal and Child Health**.

In Turkey, women's fertility is linked to education, employment and freedom to choose a husband, see **Middle East**.

Sterilization
Female sterilization in Nepal: a comparison of two types of service delivery, see **Asia**.

Teenagers
Gender differences in factors influencing first intercourse among urban students in Chile, see **Sexual Behavior**.

Sexual behavior and attitudes among unmarried urban youths in Guinea, see **Africa**.

Teenage contraceptive needs in urban South Africa: a case study, see **Africa**.

Author Index

Abdel-Tawab N, see Huntington D.

Adewole IF, see Henshaw SK.

Agadjanian V, "Quasi-legal" abortion services in a sub-Saharan setting: users' profile and motivations, p. 111.

Agha S, Sexual activity and condom use in Lusaka, Zambia, p. 32.

Ahmed MK, Rahman M and van Ginneken J, Induced abortion in Matlab, Bangladesh: trends and determinants, p. 128.

Attafuah JD, see Steiner MJ.

Bankole A, see Westoff CF.

Bankole A and Singh S, Couples' fertility and contraceptive decision-making in developing countries: hearing the man's voice, p. 15; and Correction, p. 138.

Bankole A, Singh S and Haas T, Reasons why women have induced abortions: evidence from 27 countries, p. 117.

Bishop A, see Trinh HV.

Carlson E and Omori M, Fertility regulation in a declining state socialist economy: Bulgaria, 1976–1995, Research Note, p. 184.

Cuca YP, see Henshaw SK.

de Pinho H, see Guttmacher S.

Doumbia F, see Katz KR.

Friedman M, see Thapa S.

Görgen R, Yansané ML, Marx M and Millimounou D, Sexual behavior and attitudes among unmarried urban youths in Guinea, p. 65.

Graham MJ, Larsen U and Xu X, Son preference in Anhui Province, China, p. 72.

Guttmacher S, Kapadia F, Naude JTW and de Pinho H, Abortion reform in South Africa: a case study of the 1996 Choice on Termination of Pregnancy Act, Special Report, p. 191.

Haas T, see Bankole A.

Hardee K, Janowitz B, Stanback J and Villinski MT, What have we learned from studying changes in service guidelines and practices? Comment, p. 84.

Hassan EO, see Huntington D.

Henshaw SK, Singh S, Oye-Adeniran BA, Adewole IF, Iwere N and Cuca YP, The incidence of induced abortion in Nigeria, p. 156.

Henshaw SK, see Rahman A.

Holt BY, see Potts M.

Hossain Z, see Schuler SR.

Hull TH, The challenge of contraceptive implant removals in East Nusa Tenggara, Indonesia, Comment, p. 176.

Huntington D, Nawar L, Hassan EO, Youssef H and Abdel-Tawab N, The postabortion caseload in Egyptian hospitals: a descriptive study, p. 25.

Iwere N, see Henshaw SK.

Janowitz B, see Hardee K.

Kané F, see Katz KR.

Kapadia F, see Guttmacher S.

Katz KR, West CG, Doumbia F and Kané F, Increasing access to family planning services in rural Mali through community-based distribution, p. 104.

Katzive L, see Rahman A.

Kim YM, Kols A and Mucheke S, Informed choice and decision-making in family planning counseling in Kenya, p. 4.

Kols A, see Kim YM.

Kun KE, Vaginal drying agents and HIV transmission, Viewpoint, p. 93.

Landry E, see Wegner MN.

Larsen U, see Graham MJ.

Luengo-Charath X, see Murray NJ.

Luong XH, see Trinh HV.

Marx M, see Görgen R.

Measham D, see Stein K.

Mfono Z, Teenage contraceptive needs in urban South Africa: a case study, Comment, p. 180.

Millimounou D, see Görgen R.

Mucheke S, see Kim, YM.

Murray NJ, Zabin LS, Toledo-Dreves V and Luengo-Charath X, Gender differences in factors influencing first intercourse among urban students in Chile, p. 139.

Naude JTW, see Guttmacher S.

Nawar L, see Huntington D.

Nguyen TI, see Trinh HV.

Nutley T, see Steiner MJ.

Omori M, see Carlson E.

Oye-Adeniran BA, see Henshaw SK.

Potts M, Rooks J and Holt BY, How to improve family planning and save lives using a stage-of-life approach, Viewpoint, p. 195.

Radovanovic Z, see Shah NM.

Rahman A, Katzive L and Henshaw SK, A global review of laws on induced abortion, 1985–1997, p. 56.

Rahman M, see Ahmed MK.

Rooks J, see Potts M.

Schuler SR and Hossain Z, Family planning clinics through women's eyes and voices: a case study from rural Bangladesh, p. 170.

Shah MA, see Shah NM.

Shah NM, Shah MA and Radovanovic Z, Patterns of desired fertility and contraceptive use in Kuwait, p. 133.

Singh S, see Bankole A; see also Henshaw SK.

Stanback J, see Hardee K; see also Steiner MJ.

Stein K, Measham D and Winikoff B, The quality of family planning services for breastfeeding women in Senegal, Research Note, p. 188.

Steiner MJ, Attafuah JD, Stanback J and Nutley T, Where have all the vaginal foaming tablets gone? program statistics and user dynamics in Ghana, Research Note, p. 91.

Thapa S and Friedman M, Female sterilization in Nepal: a comparison of two types of service delivery, p. 78.

Toledo-Dreves V, see Murray NJ.

Tran DC, see Trinh HV.

Trinh HV, Bishop A, Vuong TH, Luong XH, Tran DC and Nguyen TI, The potential impact of introducing pregnancy testing into menstrual regulation services in Vietnam, p. 165.

Tzanis J, see Wegner MN.

van Ginneken J, see Ahmed MK.

Villinski MT, see Hardee K.

Vuong TH, see Trinh HV.

Wegner MN, Landry E, Wilkinson D and Tzanis J, Men as partners in reproductive health: from issues to action, Special Report, p. 38.

West CG, see Katz KR.

Westoff CF and Bankole A, The time dynamics of unmet need: an example from Morocco, p. 12.

Wilkinson D, see Wegner MN.

Winikoff B, see Stein K.

Xu X, see Graham MJ.

Yansané ML, see Görgen R.

Youssef H, see Huntington D.

Zabin LS, see Murray NJ